

On the Entropy of Minimally Coupled and Non-minimally Coupled Gravities

Jeongwon Ho*, Yongduk Kim and Young-Jai Park[†]

Department of Physics and Basic Science Research Institute
Sogang University, C.P.O.Box 1142, Seoul 100-611, Korea

Abstract

We investigate whether the gravitational thermodynamic properties of the scalar-tensor theory of gravity are affected by the conformal transformation or not. As an explicit example, we consider an electrically charged static spherical black hole in the 4-dimensional low energy effective theory of bosonic string.

*E-mail address : jwho@physics.sogang.ac.kr

[†]E-mail address : yjpark@physics.sogang. ac.kr

I. Introduction

The conformal transformation of the action has been studied for various reasons. The usual actions in massless field theories are known to have conformal (Weyl) invariance, and ambiguities do not appear at the classical level [1]. Especially, in the scalar-tensor theories of gravity, one frequently transforms the action to an Einstein-Hilbert form for convenience of calculation and so on. This is based on the fact that the conformal transformation does not change the physics of the system, but just shrinks or stretches the manifold [2]. However, we note that in many cases the surface terms in the gravitational action play an important role in the action principle, and these terms are affected by a conformal transformation.

The purpose of this paper is to study in detail the effect of a conformal transformation on the scalar-tensor theory of gravity. In particular, we will focus our attention on the total energy and the entropy of a gravitational system. These will be derived from both the action describing the system and the conformally transformed one. And we will compare them. In the next section, we carry out the formulation of the total energy of the spatially bounded gravitational system. It is derived from both the nonminimally coupled action and the conformally transformed action (the minimal action) by using the gravitational Hamiltonians. In section III, the 4-dimensional low energy effective theory of bosonic string is investigated as an explicit example of the formulation. And the total energies observed in asymptotic region ($r \rightarrow \infty$) are calculated in both cases. In section IV, we compute the black hole entropy in the 4-dimensional low energy effective theory of bosonic string. Section V contains the conclusion.

II. Formulation of the total energies

Consider a manifold $(M, g_{\mu\nu})$ with the boundary ∂M which consists of initial and final spacelike surfaces ($\Sigma_{t'}$ and $\Sigma_{t''}$, respectively) and a surface near infinity (timelike three-surface Σ^∞). We assume that the spacelike surfaces Σ_t is orthogonal to Σ^∞ at the two-dimensional spacelike boundary S_t^∞ of Σ_t . The orthogonality means that on the boundary Σ^∞ , the timelike unit normal n^μ to Σ_t and the spacelike unit normal r^μ to Σ^∞ satisfy the

relation $(n \cdot r)|_{\Sigma^\infty} = 0$. In other words, the product of S_t^∞ with segments of timelike world lines orthogonal to Σ_t at S_t^∞ is the three-boundary Σ^∞ . The extrinsic curvatures of Σ_t and Σ^∞ as embedded in M are denoted $K_{ij} = h_i^k \nabla_k n_j$ and $\Theta_{ij} = \gamma_i^k \nabla_k r_j$, respectively. The extrinsic curvature of the two-dimensional spacelike boundary S_t^∞ as embedded in Σ_t is $k_{ab} = \sigma_a^k D_k r_b$, where h_{ij} , γ_{ij} and σ_{ab} are the induced metrics of Σ_t , Σ^∞ and S_t^∞ , respectively. The symbols ∇ and D denote covariant differentiations with respect to the metrics $g_{\mu\nu}$ and h_{ij} , respectively.

In the spacetime M , if a scalar field Φ is nonminimally coupled to scalar curvature, the action describing the system can be written as

$$I = \int_M d^4x \sqrt{-g} \left(\frac{1}{16\pi} \Phi R + \mathcal{L}_m \right) + \frac{1}{8\pi} \int_{\Sigma_t} d^3x \sqrt{h} \Phi K + \frac{1}{8\pi} \int_{\Sigma^\infty} d^3x \sqrt{-\gamma} \Phi \Theta, \quad (1)$$

where $\mathcal{L}_m$ is the matter Lagrangian density which is assumed to involve at most the first derivative and does not contain the gauge field. The surface term is required so that the action yields the correct equations of motion subject only to the condition that the boundary variables, which are induced three metrics and matter fields on the boundary, are held fixed. Now we can obtain a new action of the Einstein-Hilbert form through an appropriate conformal transformation from the action (1).

$$\tilde{I} = \int_M d^4x \sqrt{-\tilde{g}} \left(\frac{1}{16\pi} \tilde{R} + \tilde{\mathcal{L}}_m \right) + \frac{1}{8\pi} \int_{\Sigma_t} d^3x \sqrt{\tilde{h}} \tilde{K} + \frac{1}{8\pi} \int_{\Sigma^\infty} d^3x \sqrt{-\tilde{\gamma}} \tilde{\Theta}, \quad (2)$$

where $g_{\mu\nu} = \Phi^{-1}(x) \tilde{g}_{\mu\nu}$.

Recently, Hawking and Horowitz have proposed the general form of the total energy for the spacetime with noncompact geometry as well as compact one [3]. They have shown that the boundary terms in the Hamiltonian come directly from the boundary terms in the action rather than being put ‘by hand’. If the static slices are labeled as $N_0 = N$ on Σ^∞ , the physical Hamiltonian derived from the action (2) is then

$$\begin{aligned} \tilde{H}_p &\equiv \tilde{H} - \tilde{H}_0 \\ &= \int_{\Sigma_t} (\tilde{N} \tilde{\mathcal{H}} + \tilde{N}^i \tilde{\mathcal{H}}_i) - \int_{S_t^\infty} \left(\frac{1}{8\pi} \tilde{N} (\tilde{k} - \tilde{k}_0) - 2 \tilde{N}^\mu \tilde{p}_{\mu\nu} \tilde{r}^\nu / \sqrt{\tilde{h}} \right), \end{aligned}$$

where ‘0’ denotes the reference background, and the following relation is used;

$$R = {}^3R + K_{\mu\nu}K^{\mu\nu} - K^2 + 2\nabla_\mu(n^\mu\nabla_\nu n^\nu) - 2\nabla_\nu(n^\mu\nabla_\mu n^\nu). \quad (3)$$

Given a solution, its total energy associated with the time translation $\tilde{t}^\mu$ is

$$\tilde{E} = - \int_{S_t^\infty} d^2x \sqrt{\tilde{\sigma}} \left[\frac{1}{8\pi} \tilde{N} \tilde{k} - 2\tilde{N}^\mu \tilde{p}_{\mu\nu} \tilde{r}^\nu / \sqrt{\tilde{h}} \right]_0^{cl}, \quad (4)$$

where $]_0^{cl}$ denotes evaluation at the classical solution minus evaluation for the chosen reference background space.

Now we generalize their procedure to the nonminimally coupled action (1). In this case, the scalar field Φ is multiplied by curvature scalar and the total derivative terms in eq.(3) can not directly contribute to the boundary terms. But, after partial integrations, we obtain the appropriate boundary terms as follows,

$$\begin{aligned} & \frac{1}{16\pi} \int_M d^4x \sqrt{-g} \Phi R \\ &= \frac{1}{16\pi} \int_M d^4x N \sqrt{h} [\Phi({}^3R + K_{\mu\nu}K^{\mu\nu} - K^2) - 4n^\mu \partial_\mu \Phi K - 2D_\mu D^\mu \Phi] \\ & \quad - \frac{1}{8\pi} \int_{\Sigma_t} d^3x \sqrt{h} \Phi K + \frac{1}{8\pi} \int_{\Sigma^\infty} d^3x \sqrt{-\gamma} [\Phi n^\mu n^\nu \nabla_\nu r_\mu + r^\mu \partial_\mu \Phi]. \end{aligned} \quad (5)$$

Substituting eq.(5) to the action (1), we can rewrite the action as following

$$\begin{aligned} I &= \frac{1}{16\pi} \int_M d^4x N \sqrt{h} [\Phi({}^3R + K_{\mu\nu}K^{\mu\nu} - K^2) - 4n^\mu \partial_\mu \Phi K - 2D_\mu D^\mu \Phi + 16\pi \mathcal{L}_m] \\ & \quad + \frac{1}{8\pi} \int_{\Sigma^\infty} d^3x N \sqrt{\sigma} (\Phi k + r^\mu \partial_\mu \Phi). \end{aligned} \quad (6)$$

Note that a new boundary term $\int_{\Sigma^\infty} r^\mu \partial_\mu \Phi$ appears in eq.(6). As a result, the total energy derived from the action (1) is

$$E = - \int_{S_t^\infty} d^2x \sqrt{\sigma} \left[\frac{1}{8\pi} N (\Phi k + r^\mu \partial_\mu \Phi) - 2N^\mu p_{\mu\nu} r^\nu / \sqrt{h} \right]_0^{cl}. \quad (7)$$

Comparing eq.(4) and eq.(7), one sees that the expressions of the total energy, which are computed from both the minimal and the nonminimal actions, are not equal to each other. However, when $\Phi = 1$, these become the same quantities. Note that the difference between $\tilde{E}$ and E is mainly due to the projection to the two-dimensional boundary of the gradient of

the scalar field. In other words, the term $(\Phi k + r^\mu \partial_\mu \Phi)$ seems to play a role of the effective extrinsic curvature associated with the boundary S_t^∞ in the nonminimal case.

III. The 4-dimensional low energy effective theory of bosonic string

Now we shall apply the results in the previous section to the 4-dimensional low energy effective theory of bosonic string as an explicit example. The action of the theory is

$$I = \int_M d^4x \sqrt{-g} e^{-2\phi} \left[\frac{1}{16\pi} R + \frac{1}{4\pi} \partial_\mu \phi \partial^\mu \phi - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} \right] + \frac{1}{8\pi} \int_{\Sigma_t} d^3x \sqrt{h} e^{-2\phi} K + \frac{1}{8\pi} \int_{\Sigma^\infty} d^3x \sqrt{-\gamma} e^{-2\phi} \Theta, \quad (8)$$

where ϕ is the dilaton field, and F is the field strength of the gauge field A . Through a conformal transformation $g_{\mu\nu} = e^{2\phi} \tilde{g}_{\mu\nu}$, we obtain a new action of the Einstein-Hilbert form as follows

$$\tilde{I} = \int_M d^4x \sqrt{-\tilde{g}} \left[\frac{\tilde{R}}{16\pi} - \frac{1}{8\pi} \partial_\mu \phi \partial^\mu \phi - \frac{1}{4} e^{-2\phi} F_{\mu\nu} F^{\mu\nu} \right] + \frac{1}{8\pi} \int_{\Sigma_t} d^3x \sqrt{\tilde{h}} \tilde{K} + \frac{1}{8\pi} \int_{\Sigma^\infty} d^3x \sqrt{-\tilde{\gamma}} \tilde{\Theta}, \quad (9)$$

where the raising and lowering of indices are carried out by the new metric $\tilde{g}_{\mu\nu}$. Varying the above two actions and solving the Einstein's equations, we have obtained the electrically charged static spherical black hole solutions as follows

$$ds^2 = -\left(1 - \frac{r_H}{r}\right) \left(1 + \frac{\alpha}{r}\right)^{-2} dt^2 + \left(1 - \frac{r_H}{r}\right)^{-1} dr^2 + r^2 d\Omega_2^2, \quad (10)$$

and

$$d\tilde{s}^2 = -\left(1 - \frac{r_H}{r}\right) \left(1 + \frac{\alpha}{r}\right)^{-1} dt^2 + \left(1 - \frac{r_H}{r}\right)^{-1} \left(1 + \frac{\alpha}{r}\right) dr^2 + r^2 \left(1 + \frac{\alpha}{r}\right) d\Omega_2^2, \quad (11)$$

respectively, where $\alpha \equiv Q^2/4\pi M$ and $r_H \equiv 2M(1 - Q^2/8\pi M^2)$ [4]. Here, Q is the electric charge and M is the ADM mass. The corresponding dilaton and gauge fields are given by

$$e^{-2\phi} = \left(1 + \frac{\alpha}{r}\right),$$

$$A_\mu = \left(\frac{Q}{4\pi r} \left(1 + \frac{\alpha}{r}\right)^{-1} - \Phi_H, 0, 0, 0 \right), \quad (12)$$

respectively, where $\Phi_H \equiv Q/8\pi M$ is the gauge fixing term chosen on the condition of regularizing the gauge field at horizon $r = r_H$.

Since both actions (8) and (9) contain the gauge field, we have another surface term $\int_{s_t^\infty} A_t p_\mu^A r^\mu$ in the process of deriving the Hamiltonian, where p_A^μ is the momenta conjugate to the gauge field A_μ . Thus the total energy for the minimal case (4) becomes [5]

$$\tilde{E} = - \int_{s_t^\infty} d^2x \sqrt{\tilde{\sigma}} \left[\frac{1}{8\pi} \tilde{N} \tilde{k} - 2 \tilde{N}^\mu \tilde{p}_{\mu\nu} \tilde{r}^\nu / \sqrt{\tilde{h}} - A_t \tilde{p}_\mu^A \tilde{r}^\mu / \sqrt{\tilde{h}} \right]_0^{cl}. \quad (13)$$

On the other hand, the total energy for the nonminimal case (7) becomes

$$E = - \int_{s_t^\infty} d^2x \sqrt{\sigma} \left[\frac{1}{8\pi} N (\Phi k + r^\mu \partial_\mu \Phi) - 2 N^\mu p_{\mu\nu} r^\nu / \sqrt{h} - A_t p_\mu^A r^\mu / \sqrt{h} \right]_0^{cl}, \quad (14)$$

where $\Phi \equiv e^{-2\phi}$.

Substituting eqs. (10 - 12) to (13) and (14), we obtain the total energies observed in asymptotic region of the black hole as

$$\tilde{E} = M - Q\Phi_H + \frac{\alpha}{2} \quad \text{and} \quad E = M - Q\Phi_H - \alpha, \quad (15)$$

respectively [6]. Here, we set $N = N_0 = 1$, $\tilde{N} = \tilde{N}_0 = 1$. Note that this restriction gives the value of the Hamiltonian, which generates unit time translations. And we have required that the reference background space is a static solution to the field equations ($p^{\mu\nu}|_0$, $p_\Phi|_0$, $p_A^\mu|_0$, and the constraints vanishes), and $\Phi|_0 = 1$.

In eq.(15), we can see that $\tilde{E}$ differs from the usual total energy of the grand canonical ensemble, $M - Q\Phi_H$, by the factor $\alpha/2$. This is due to the dilaton charge $\mathcal{D}$ observed in infinite region

$$\mathcal{D} \equiv \frac{1}{4\pi} \int_{\Sigma_t^\infty} d^2x \sqrt{\tilde{\sigma}} \tilde{r}^\mu \nabla_\mu \phi = \frac{\alpha}{2} + \mathcal{O}(r^{-1}). \quad (16)$$

Qualitatively, this result is originated as follows: although the dilaton charge is not an independent degree of freedom in the grand canonical ensemble ($\alpha = Q^2/4\pi M$), this characterizes the black hole solution in the asymptotic region [7]. So it seems natural for the dilaton charge to be included in the total energy observed in asymptotic region. On the

other hand, since the values of $\tilde{E}$ and E differ by the factor $3\alpha/2$, we can say that the total energy observed in asymptotic region is changed by conformal transformation.

V. The black hole entropy

In this section, we shall show that the black hole entropies computed from two actions (8) and (9) are equal to each other at the semiclassical limit [8]. Calculating the entropy from the Einstein-Hilbert action (9) and the corresponding black hole solution (11), we will show that the entropies calculated via well-known three methods are equal to each other. It is important to note that we use the results in eq.(15) for the entropies of the black holes with the Gibbons-Hawking method [8].

Firstly, from the Bekenstein-Hawking relation [9,10], we obtain

$$\tilde{S}_m = \frac{1}{4}A^H = 4\pi M^2 \left(1 - \frac{Q^2}{8\pi M^2}\right). \quad (17)$$

Secondly, from the first law of black hole thermodynamics, $T_H dS = dM - \Phi_H dQ$ [11] (the electrically charged black hole in eq.(11) is characterized by $(M, Q, J = 0)$), the black hole entropy is given by

$$\tilde{S}_m = \int \frac{dM}{\tilde{T}_H} - \int \frac{\Phi_H}{\tilde{T}_H} dQ = 4\pi M^2 \left(1 - \frac{Q^2}{8\pi M^2}\right), \quad (18)$$

where the Hawking temperature $\tilde{T}_H$ is $\tilde{T}_H = T_H = \mathcal{K}/2\pi = 1/8\pi M$ and $\mathcal{K}$ is the surface gravity of the black hole.

Next, let us compute the entropy with the method suggested by Gibbons and Hawking [8]. The Euclidean action is given by

$$\begin{aligned} \tilde{I}_E[\tilde{g}, A, \phi] = & - \left[\int_Y d^4x \sqrt{-\tilde{g}} \left(\frac{\tilde{R}}{16\pi} - \frac{1}{8\pi} \partial_\mu \phi \partial^\mu \phi \right) + \frac{1}{8\pi} \int_{\partial Y} d^3x \sqrt{\tilde{\gamma}} (\tilde{\Theta} - \tilde{\Theta}_0) \right] \\ & + \frac{1}{4} \int_Y d^4x \sqrt{\tilde{g}} e^{-2\phi} F_{\mu\nu} F^{\mu\nu}, \end{aligned} \quad (19)$$

where the 4-dimensional Euclidean manifold $Y = \mathfrak{R} \times S^1 \times S^2$, and its boundary $\partial Y = S^1 \times S^2$.

Then, the gravitational partition function and the entropy are given by

$$\begin{aligned}\tilde{Z} &= \int [d\tilde{g}_{\mu\nu}][dA_\mu][d\phi] e^{-\tilde{I}[\tilde{g}, A, \phi]} \simeq e^{-\tilde{I}[\tilde{g}_c, A_c, \phi_c]}, \\ \tilde{S}_m &\simeq -\tilde{I}[\tilde{g}_c, A_c, \phi_c] + \frac{1}{\tilde{T}_H} \left(M - Q\Phi_H + \frac{\alpha}{2} \right),\end{aligned}\tag{20}$$

where the subscript ‘c’ denotes the saddle point of the Euclidean action and

$$\tilde{I}_E[\tilde{g}_c, A_c, \phi_c] = \frac{1}{2\tilde{T}_H} (M - Q\Phi_H + \alpha) + \mathcal{O}(r^{-1}).\tag{21}$$

Therefore, the black hole entropy is obtained as follows

$$\tilde{S}_m \simeq 4\pi M^2 \left(1 - \frac{Q^2}{8\pi M^2} \right).\tag{22}$$

This is the same as the results in eqs.(17) and (18) as expected. Here, it must be emphasized that we did not use $(M - Q\Phi_H)$ as the total energy of the system, but $(M - Q\Phi_H + \alpha/2)$ derived in the previous section. Note that if we substitute $(M - Q\Phi_H)$ in eq.(20), we could not have obtained the same black hole entropy with which evaluated in alternative ways.

On the other hand, for the nonminimal case we can also compute the black hole entropy via Gibbons-Hawking method. The Euclidean action is given by

$$\begin{aligned}I_E[g, A, \phi] &= - \left[\int_Y d^4x \sqrt{g} e^{-2\phi} \left(\frac{R}{16\pi} + \frac{1}{4\pi} \partial_\mu \phi \partial^\mu \phi \right) + \frac{1}{8\pi} \int_{\partial Y} d^3x \sqrt{\gamma} e^{-2\phi} (\Theta - \Theta_0) \right] \\ &\quad + \frac{1}{4} \int_Y d^4x \sqrt{g} e^{-2\phi} F_{\mu\nu} F^{\mu\nu}.\end{aligned}\tag{23}$$

In the saddle point approximation, the first integration in eq.(23) does not vanish via the equations of motion. This term gives another surface term as follows

$$- \frac{3}{16\pi} \int_{\partial Y} d^3x \sqrt{\gamma} (g_{rr})^{-1/2} \partial_r \left(1 + \frac{\alpha}{r} \right) = \frac{3\alpha}{4} \left(\frac{2\pi}{\mathcal{K}} \right) + \mathcal{O}(r^{-1}).$$

Then, we obtain the value of the action in saddle point approximation

$$I_E[g_c, A_c, \phi_c] = \frac{1}{2T_H} (M - Q\Phi_H - 2\alpha) + \mathcal{O}(r^{-1}).$$

As a result, the black hole entropy becomes

$$S_n \simeq -I[g_c, A_c, \phi_c] + \frac{1}{T_H} (M - Q\Phi_H - \alpha) \simeq 4\pi M^2 \left(1 - \frac{Q^2}{8\pi M^2} \right).\tag{24}$$

Therefore, we have shown that $S_n = \tilde{S}_m$. From this result, one recognize that the black hole entropy is not scaled by the conformal transformation up to the semiclassical limit.

VI. Conclusion

In this paper, we have explicitly examined whether the total energy and the entropy of a gravitational system, in particular a black hole, are affected by the conformal transformation or not. The total energy is computed via deriving the gravitational Hamiltonian from the action. As a result, we have shown that the total energy computed from the conformally transformed action (minimal action) is different with that computed from the original action (nonminimal action) such that the extrinsic curvature associated with the two-dimensional boundary S_t^∞ , $\tilde{k}$ in the former is replaced with the *effective extrinsic curvature* $(\Phi k + r^\mu \partial_\mu \Phi)$ in the latter.

In the 4-dimensional low energy effective theory of bosonic string, the total energies observed in asymptotic region of the electrically charged static spherical black hole are given in eq.(15). They differ by the factor $3/2\alpha$, and the factor is related with the dilaton charge $\mathcal{D}$. So, we can say that the total energy observed in asymptotic region is scaled by the conformal transformation. Also, we note that the total energies are different from the usual total energy of the grand canonical ensemble, $(M - Q\Phi_H)$. In order to understand it we have to consider two factors related each other. One is the method computing total energy. We have considered a timelike boundary, and then it is sent back to infinite region. Another is that although the dilaton charge is not the independent degree of freedom in the grand canonical ensemble, the black hole solution is characterized by the dilaton charge as well as the ADM mass and the electric charge in asymptotic region. In other words, owing to the method, the effect of the dilaton charge is contained in the total energy of the gravitational system.

In the case of the action of the Einstein-Hilbert form, the black hole entropy is calculated in the well-known three methods. And all of these three methods have yielded the same

expressions for entropy. On the process of the calculation via the Gibbons-Hawking method, it is the important point that we did not use $(M - Q\Phi_H)$ as the total energy of the black hole, but $(M - Q\Phi_H + \alpha/2)$. On the other hand, when the black hole entropy is computed from the nonminimally coupled action via the Gibbons-Hawking method, also we have used the result in eq.(15) as the total energy. As a result, we have shown that this gives the same expression for the entropy with that calculated from the Einstein-Hilbert form. Furthermore, we have obtained the same Hawking temperature in two cases. Of course, the local (Tolman redshift) temperatures, $T(r) = (-g_{tt})^{-1/2}T_H$ and $\tilde{T}(r) = (-\tilde{g}_{tt})^{-1/2}\tilde{T}_H$, are different each other.

In conclusion, the conformal transformation changes local geometry, while Hawking temperature T_H and black hole entropy S are global quantities. Therefore, the values of T_H and S are invariant under the conformal transformation [12, 13].

Acknowledgments

This work was supported by Basic Science Research Institute Program, Ministry of Education, Project No. 95-2414.

References

- [1] S. B. Treiman, R. Jackiw, B. Zumino and E. Witten, *Current Algebra and Anomalies* (Princeton University Press, Princeton, 1985).
- [2] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).
- [3] S. W. Hawking and G. T. Horowitz, "The Gravitational Hamiltonian, Action, Entropy and Surface terms", gr-qc/9501014. J. D. Brown and J. W. York, Jr., Phys. Rev. **D47**, 1407(1993).
- [4] G. W. Gibbons and K. Maeda, Nucl. Phys. **B298**, 741(1988). I. Ichinose and H.

- Yamazaki, Mod. Phys. Lett. **A4**, 1509(1989).
- [5] S. W. Hawking and S. F. Ross, “Duality between Electric and Magnetic Black Holes”, hep-th/9504019.
- [6] Compare with the result of J. Katz, D. Lynden-Bell, and W. Israel, Class. Quantum Grav. **5**, 971(1988).
- [7] D. Garfinkle, G. Horowitz and A. Strominger, Phys. Rev. D**43**, 3140(1991).
- [8] G. W. Gibbons and S. W. Hawking, Phys. Rev. D**15**, 2752(1977). For rotating black holes, J. D. Brown, E. A. Martinez and J. W. York, Jr., Phys. Rev. Lett. **66**, 2281(1991).
- [9] J. D. Bekenstein, Phys. Rev. D**7**, 2333(1973); *ibid* D**9**, 3292(1974).
- [10] S. W. Hawking, Commun. Math. Phys. **43**, 199(1975).
- [11] J. W. Bardeen, B. Carter and S. W. Hawking, Comm. Math. Phys. **31**, 161(1973).
- [12] R. Kallosh, T. Ortín and A. Peet, Phys. Rev. D**47**, 5400(1993).
- [13] K. C. K. Chan, J. D. E. Creighton and R. B. Mann, Phys. Rev. D**54**, 3892(1996).